



# New record of the Jaguar, *Panthera onca* (Linnaeus, 1758) (Felidae), from a mosaic of Atlantic Forest in the Paraná state, Brazil

Diogo C. Casanova<sup>1</sup> & Rodrigo Bernardo

Casa da Floresta Ambiental SS. Av. Joaquinha Morganti, 289. CEP 13415-030. Piracicaba, São Paulo, Brazil

<sup>1</sup> Corresponding author. E-mail: [diogo@casadafloresta.com.br](mailto:diogo@casadafloresta.com.br)

**Abstract:** The largest felid in the Americas, the Jaguar (*Panthera onca*) has had its geographical distribution reduced to approximately half the original range throughout the years, and most of its remaining distribution is in Brazilian territory. Deforestation is considered the main threat to the species, especially in the Atlantic Rainforest, which is currently highly fragmented and modified. This article provides a new record of Jaguar in Paraná state, in an area occupied by a mosaic formed by native vegetation and silviculture, and represents a new recent locality of the species for the biome in which it is classified as Critically Endangered of extinction.

**Key words:** Carnivora; Monte Alegre Farm; silvicultural landscape; Telêmaco Borba

Considered the largest felid species in the Americas, the Jaguar, *Panthera onca* (Linnaeus, 1758), is the only species of the genus found in the New World (MACDONALD et al. 2010; MARTINS 2015). Originally, this species range extended from northern Argentina to North America, covering a large portion of southern United States (SANDERSON et al. 2002; ZELLER 2007). Currently, its range is only half of the original, and approximately 50% of this current range is within Brazil (ZELLER 2007). Additionally, this species is locally extinct in several countries, such as Uruguay and El Salvador (CASO et al. 2008).

The main threats for this species are anthropic modification of landscapes (RIPPLE et al. 2014), resulting in habitat loss and fragmentation (MORATO et al. 2013; VILLARD & METZGER 2014), reduction in prey abundance (RYALL & FAHRIG 2006; AHUMADA et al. 2011), and increase of hunting (WHITE & LOWE 2008; INSKIP & ZIMMERMANN 2009), motivated mainly by conflicts generated by predation of livestock (CASO et al. 2008; PALMEIRA et al. 2008; PAVILO et al. 2008).

The Jaguar is extant in all Brazilian biomes, with the exception of the Pampa (TORRES et al. 2008; NASCIMENTO

& CAMPOS 2011), and its presence and abundance depend on factors such as prey availability, presence of water bodies, and well-preserved native vegetation (EISENBERG & REDFORD 1999; SUNQUIST & SUNQUIST 2002; ARROYO-ARCE et al. 2014). In the Brazilian Atlantic rainforest, where fragmentation and modification of the landscape have been intense, it is estimated that in the past 15 years the population of this felid has been reduced by at least 80% (MORATO et al. 2013). Currently, it is believed that the species is divided into eight subpopulations within the Atlantic Rainforest (BEISIEGEL et al. 2012) and the total population size does not exceed 300 individuals (PAVILO et al. 2016).

Considered as one of Earth's 34 biodiversity hotspots (MYERS et al. 2000; MITTERMEIER et al. 2004), the Atlantic Rainforest historically extended north to south along the Brazilian coast, inland reaching as far as Argentina and Paraguay, and covering an area of 1,315,460 km<sup>2</sup> (IBGE 2008). Deforestation represents the main threat for this biome, which is currently reduced to approximately 12% of its original size (RIBEIRO et al. 2009). This drastic reduction in the available habitat, associated with constant threats to the survival of the species, account for the categorization of the Jaguar as Critically Endangered in this biome (BEISIEGEL et al. 2012; PAVILO et al 2016).

In Paraná state, southern Brazil, the range of the Jaguar is divided in three regions, as follows: 1) Foz do Iguaçu Region (Green Corridor), on the border between Brazil and Misiones, Argentina; 2) Extreme Northwest, between the states of São Paulo, Paraná, and Mato Grosso do Sul; 3) Serra do Mar / South, in the Serra do Mar range (between the states of São Paulo and Paraná) and municipalities in the southern part of Paraná state (LIMA et al. 2013). The Jaguar subpopulation of Paraná is one of six along the Brazilian Atlantic coast (HAAG et al 2010; BEISIEGEL et al. 2012).

The pelt of this species is yellowish, with dark rosetted spots, and differently from other Neotropical felids, the



**Figure 1.** Captures of the camera trap filming of the Jaguar, *Panthera onca*, on Monte Alegre Farm, Telêmaco Borba, Paraná state, in 2016.

rosettes are larger, fewer, and often having one or more black dots inside (KITCHENER 1991; SUNQUIST & SUNQUIST 2002). The Jaguar has a muscular body, broad chest, and relatively shortlimbs, with the rounded head large compared to the rest of the body (SEYMOUR 1989) (Figure 1). Moreover, it can be easily diagnosed from the only other large felid in the region, Puma (*Puma concolor* (Linnaeus, 1771)), which is smaller, with a relatively longer tail, and self-colored (CURRIER 1983).

Monte Alegre Farm (24°12'42"S, 050°33'26"W), owned by Klabin S.A., is a 269,700-ha sylvicultural farm located in the municipality of Telêmaco Borba, in central Paraná state. The local landscape forms a mosaic of native vegetation (110,000 ha) and silviculture (134,800 ha), with the latter divided among cultivated areas of *Eucalyptus* sp., *Pinus* sp., and *Araucaria angustifolia*. This forested area of the whole Monte Alegre Farm (Figure 2), although partially occupied by exotic species, represents an important remnant in Paraná, since it represents in terms of forest coverage in the state a relatively large portion of the preserved native vegetation, formed by an area of ecological tension between the Mixed Ombrophilous Forest and the Cerrado (REIS et al. 2006; IBGE 2012).


Inside the study area, medium-sized and large mammals have been monitored since 2011, through campaigns held twice yearly. The plantation matrix and four main fragments with native forest were monitored, with two areas sampled per campaign (the average size of the fragments is 1,580.4 ha). In each campaign, a total of 18 camera traps were used, nine in each fragment, and programmed to operate uninterruptedly for five days. This resulted in a sampling effort of 2,160 hours per campaign, totaling 12,960 hours throughout the six years monitored.

Monitoring was conducted through the use of camera traps, one of the most commonly used methods for sampling medium-sized and large mammals (LYRA-JORGE et al. 2014)


In 2016, with the use of camera traps, one adult Jaguar (unidentified sex) was recorded (Figure 1) moving along a road that crosses a field of *Eucalyptus* sp. Here, one side of the road had recent been harvested of *Eucalyptus* and was clear; the areas adjacent were fragments of native vegetation and fields of *Pinus* sp. This sighting is located 125 km west of the nearest record, a documented sighting in 1994 in the municipality of Palmeira (MEDELLIN et al. 2001) (Figure 3). Until now, the Jaguar was thought to be probably extirpated from Monte Alegre Farm because no reliable documented sightings had been obtained for over 50 years (REIS et al. 2006).

Among the factors that explain the persistence of wild mammals in modified landscapes, the quality and permeability of the surrounding matrix is considered to be one of the most important (UMETSU et al. 2008; SANTOS-FILHO et al. 2012). Compared with other land uses, such as extensive agricultural activities, silviculture may have less impact on local wildlife. The vertical structure in areas of silviculture resembles natural vegetation and there is a possibility that the understory has native vegetation (COELHO et al. 2014). Together, these become a determining matrix for the success of viable populations in areas under anthropic pressures (BOYLE & SMITH 2010).

In addition to the characteristics of the matrix, the presence or absence of species relies on the availability of resources (TRAINOR et al. 2014). In the study area, there several potential prey species for the Jaguar (GARLA et al. 2001; CULLEN JR et al. 2013). Collared Peccary (*Pecari tajacu*),



**Figure 2.** Location of Monte Alegre Farm (study area) in Paraná state, Brazil, and Atlantic Forest remnants in the region (RIBEIRO et al. 2009).



**Figure 3.** Map of the records of the jaguar *Panthera onca*, indicating priority areas for connectivity between the subpopulations of the species (pink) (Figure reproduced with authorization from PANTHERA 2016). Records according to MARES et al. (1981), EISENBERG (1989), REDFORD & EISENBERG (1992), CALOURO (1999), EISENBERG & REDFORD (1999), ORTEGA-HUERTA et al. (1999), NUÑES et al. (2000), GARLA et al. (2001), MEDELLIN et al. (2001), RODRIGUES et al. (2002), POLISAR et al. (2003), SCOGNAMILLO et al. (2003), MAFFEI et al. (2004), SILVER et al. (2004), DURDEN et al. (2006), HAAG et al. (2010), MORAES (2012), LIMA et al. (2013), and PAVIOLÓ et al. (2016).

White-lipped Peccary (*Tayassu pecari*), several armadillos (*Cabassous tatuay*, *Dasyurus spp.*, *Euphractus sexcinctus*), Coati (*Nasua nasua*), and Crab-eating Raccoon (*Procyon cancrivorus*) have been documented (REIS et al. 2006).

Among all recent records of the Jaguar in Paraná, this study is the first to report this species in a locality without a known subpopulation. It is believed that the observed individual belongs to the Serra do Mar subpopulation, given the proximity of this record to others in eastern Paraná, and is using the farm as a corridor. There is not much information about the current situation of the species in the coastal region of the Atlantic Forest in Paraná state (BEISIEGEL et al. 2012). A third of all scientific production on the ecology of the Jaguar in Brazil is concentrated in the Pantanal region (CAVALCANTI et al. 2010; SOLLmann 2010).

Additionally, Monte Alegre Farm is located near an area identified by RABINOWITZ et al. (pers. comm.) as important for the connectivity between remnant subpopulations of the Jaguar (Figure 3). The new record from Monte Alegre Farm suggests that silviculture matrices are permeable for large carnivores (CAMPOS 2009; LYRA-JORGE et al. 2011) and corroborates the findings by LIMA et al. (2013) in Mato Grosso do Sul state. Consequently, future studies are needed on the Jaguar in Paraná, as well as the permeability of the silviculture matrix, because the Monte Alegre Farm sighting was from outside the area thought by PAVIOLI et al. (2016) as being potentially occupied in the present day.

The native vegetation of Paraná has suffered greatly and is now highly fragmented and/or modified. Native vegetation currently covers 3.7 million ha, which represents only 18.51% of the original area (ACCIOLY 2013). The establishment of ecological corridors in Paraná is needed for Jaguar conservation. Also needed is a better understanding of how land use and occupation influences the ecological dynamics of this species, and especially in terms of silviculture, which with 880,000 ha occupied (*Eucalyptus* and *Pinus* (IBA 2015) together represent 4.41% of the state's territory) is prevalent in Paraná.

## ACKNOWLEDGEMENTS

The authors thank Vagner de Araújo Gabriel, Elson Fernandes de Lima, Raisa Reis de Paula Rodarte, and the two anonymous referees for the comments and suggestions. Thanks are also due to the Panthera Foundation for information provided. Thanks are also due to Fernando Igor de Godoy and Marcelo Fabrício Lino for their support and company during fieldwork. This study was supported by Klabin S.A. as part of the terrestrial fauna monitoring conducted by Casa da Floresta Ambiental SS.

## LITERATURE CITED

- ACCIOLY, P. 2013. Mapeamento dos remanescentes vegetais arbóreos do estado do Paraná e elaboração de um sistema de informações geográficas para fins de análise ambiental do estado [Thesis] Curitiba: Universidade Federal do Paraná. 127 pp.
- AHUMADA, J.A., C.E.F. SILVA, K. GAJAPERSAD, C. HALLAM, J. HURTADO, E. MARTIN, A. MCWILLIAM, B. MUGERWA, T. O'BRIEN, F. ROVERO, D. SHEL, W.R. SPIRONELLO, N. WINARNI & S.J. ANDELMAN. 2011. Community structure and diversity of tropical forest mammals: data from a global camera trap network. Philosophical transactions of the Royal Society of London 366: 2703–2711. doi: [10.1098/rstb.2011.0115](https://doi.org/10.1098/rstb.2011.0115)
- ARROYO-ARCE, S., J. GUILDER & R. SALON-PÉREZ. 2014. Habitat features influencing Jaguar *Panthera onca* (Carnivora: Felidae) occupancy in Tortuguero National Park, Costa Rica. Revista de Biología Tropical. 62(4): 1449–1458. doi: [10.15517/rbt.v63i3.16537](https://doi.org/10.15517/rbt.v63i3.16537)
- BEISIEGEL, B.M., D.A. SANA, & E.A. MORAES JR. 2012. The Jaguar in the Atlantic Forest. Cat News Special Issue 7: 14–18. [http://www.catsg.org/fileadmin/filesharing/3.Conservation\\_Center/3.2\\_Status\\_Reports/Jaguar/Beisiegel\\_et\\_al\\_2012\\_The\\_jaguar\\_in\\_the\\_Atlantic\\_Forest.pdf](http://www.catsg.org/fileadmin/filesharing/3.Conservation_Center/3.2_Status_Reports/Jaguar/Beisiegel_et_al_2012_The_jaguar_in_the_Atlantic_Forest.pdf)
- BOYLE, S.A. & A.T. SMITH. 2010. Can landscape and species characteristics predict primate presence in forest fragments in the Brazilian Amazon? Biological Conservation 143: 1134–1143. doi: [10.1016/j.biocon.2010.02.008](https://doi.org/10.1016/j.biocon.2010.02.008)
- CALOURO, A.M. 1999. Riqueza de mamíferos de grande e médio porte do Parque Nacional da Serra do Divisor (Acre, Brasil). Revista Brasileira de Zoologia 16: 195–213. doi: [10.1590/S0101-8175199900600020](https://doi.org/10.1590/S0101-8175199900600020)
- CAMPOS, C. B. 2009. Dieta de carnívoros e uso do espaço por mamíferos de médio e grande porte em áreas de silvicultura do estado de São Paulo, Brasil [thesis]. Piracicaba: Universidade de São Paulo. 137 pp.
- CARBONE, C., S. CHRISTIE, K. CONFORTI, T. COULSON, N. FRANKLIN & J.R.I. GINSBERG. 2001. The use of photographic rates to estimate densities of tigers and other cryptic mammals. Animal Conservation 4: 75–79. doi: [10.1017/S1367943001001081](https://doi.org/10.1017/S1367943001001081)
- CASO, A., C. LOPEZ-GONZALEZ, E. PAYAN, E. EIZIRIK, T. DE OLIVEIRA, R. LEITE-PITMAN, M. KELLY & C. VALDERRAMA. 2008. *Panthera onca*, in: IUCN. The Red List of threatened species. Version 2016.1. Accessed at <http://www.iucnredlist.org>, 12 September 2016.
- CAVALCANTI, S.M.C., S. MARCHINI, A. ZIMMERMANN, E.M. GESE & D.W. MACDONALD. 2010. Jaguars, livestock, and people in Brazil: realities and perceptions behind the conflict, pp. 2383402, in: D.W. MACDONALD & A.L. LOVERIDGE (eds.). Biology and conservation of wild felids. New York: Oxford University Press.
- COELHO, M., L. JUEN & A.C.M. OLIVEIRA. 2014. The role of remnants of Amazon savanna for the conservation of Neotropical mammal communities in eucalyptus plantations. Biodiversity Conservation 23(13): 3171–3184. doi: [10.1007/s10531-014-0772-9](https://doi.org/10.1007/s10531-014-0772-9)
- CULLEN JR., L., D.A. SANA, F. LIMA, K.C. ABREU, A. UEZU. 2013. Selection of habitat by the Jaguar, *Panthera onca* (Carnivora: Felidae), in the upper Paraná River, Brazil. Zoologia 30(4): 379–387. doi: [10.1590/S1984-46702013000400003](https://doi.org/10.1590/S1984-46702013000400003)
- CURRIER, M.J.P. 1983. Mammalian species: *Felis concolor*. The American Society of Mammalogists 200: 1–7. <http://www.science.smith.edu/msi/pdf/i0076-3519-200-01-0001.pdf>
- DURDEN, L.A., M.W. CUNNINGHAM, R. MCBRIDE & B. FERREE. 2006. Ectoparasites of free-ranging Pumas and Jaguars in the Paraguayan Chaco. Veterinary Parasitology 137: 189–193. doi: [10.1016/j.vetpar.2006.01.001](https://doi.org/10.1016/j.vetpar.2006.01.001)
- EISENBERG, J.F. 1989. Mammals of the Neotropics: the northern Neotropics. Vol 2. Chicago: University of Chicago Press. 449 pp.
- EISENBEG, J.F. & K.H. REDFORD. 1999. Mammals of the Neotropics: the central Neotropics. Vol. 3. Chicago: University of Chicago Press. 609 pp.
- GARLA, R.C., E.F. SETZ & N. GOBBI. 2001. Jaguar (*Panthera onca*) food habits in Atlantic Rain Forest of southeastern Brazil. Biotropica 33(4): 691–696. doi: [10.1111/j.17447429.2001.tb00226.x](https://doi.org/10.1111/j.17447429.2001.tb00226.x)
- HAAG, T., A.S. SANTOS, D.A. SANA, R.G. MORATO, R. CULLEN JR., P.G. CRAWSHAW JR., C. DE ANGELO, M. S. DI BITETTI, F.M. SALZANO

- & E. EIZIRIK. 2010. The effect of habitat fragmentation on the genetic structure of a top predator: loss of diversity and high differentiation among remnant populations of Atlantic Forest Jaguars (*Panthera onca*). *Molecular Ecology* 19(22): 4906–4921. doi: [10.1111/j.1365-294X.2010.04856.x](https://doi.org/10.1111/j.1365-294X.2010.04856.x)
- IBA (INDÚSTRIA BRASILEIRA DE ÁRVORES). 2015. Relatório IBA. Brasília: Studio 113. 78 pp.
- IBGE (INSTITUTO BRASILEIRO DE GEOGRAFIA E ESTATÍSTICA). 2008. Mapa da área da aplicação da Lei 11.428/2006. 1:5.000.000 Rio de Janeiro: IBGE
- IBGE (INSTITUTO BRASILEIRO DE GEOGRAFIA E ESTATÍSTICA). 2012. 2nd ed. 1:5,000,000. Brasília: IBGE.
- INSKIP, C. & A. ZIMMERMANN. 2009. Human-felid conflict: a review of patterns and priorities worldwide. *Oryx Conservation Journal* 43:18–34. doi: [10.1017/S003060530899030x](https://doi.org/10.1017/S003060530899030x)
- KITCHENER, A. 1991. The natural history of the wild cats. New York: Cornell University Press. 280 pp.
- LIMA, E. F., D.H. HOMEM & P.F. COLAS-ROSAS. 2013. Mammalia, Felidae, *Panthera onca* (Linnaeus, 1758): recent records in east Mato Grosso do Sul, Brazil. *Check List* 9(1): 121–124. doi: [10.15560/9.1.121](https://doi.org/10.15560/9.1.121)
- LYRA-JORGE, M.C., C. GHELER-COSTA, C.I. PINA, L.M. ROSALINO & L.M. VERDADE. 2014. Wildlife surveys in agricultural landscapes: terrestrial medium- to large-sized mammals, pp. 133–147, in: L.M. VERDADE, M.C. LYRA-JORGE, C.I. PINA (eds). Applied ecology and human dimensions in biological conservation. Berlin/Heidelberg: Springer.
- LYRA-JORGE, M.C., G. CIOCHETI, L.R. TAMBOSI, M.C. RIBEIRO & V.R. PIVELLO. 2011. Carnivorous mammals in a mosaic landscape in southeastern Brazil: Is it possible to keep them in an agro-silvicultural landscape?; pp. 317–32, in: J. RUDAS & T. DAHLGREN (org.). Grassland biodiversity — Habitat types, ecological processes and environmental impacts. New York: Nova Science Publishers.
- MACDONALD, D.W., A. L. LOVERIDGE & K. NOWELL. 2010. Dramatis personae: an introduction to the wild felids; pp. 3–58, in: D.W. MACDONALD & A.L. LOVERIDGE (eds.). Biology and conservation of wild felids. New York: Oxford University Press.
- MAFFEI, L.; E. CUÉLLAR & A. NOSS. 2004. One thousand Jaguars (*Panthera onca*) in Bolivia's Chaco? Camera trapping in the Kaa-Iya National Park. *Journal of Zoology* 262: 295–304. doi: [10.1017/S0952836903004655](https://doi.org/10.1017/S0952836903004655)
- MARES, M.A., M.R. WILLIG, K.E. STREILEIN & E.T.E. LACHER JR. 1981. The mammals of northeastern Brazil: a preliminary assessment. *Annals of Carnegie Museum of Natural History* 50 (4): 81–137.
- MARTINS, R. 2015. Conservação de onça-parda (*Puma concolor*) e de onça-pintada (*Panthera onca*) no mosaico da Juréia-Itatins, São Paulo [Dissertation]. Santos: Universidade Santa Cecília. 124 pp.
- MEDELLIN, R.A., C. CHETKIEWICZ, A. RABINOWITZ, K.H. REDFORD, J.G. ROBINSON, E.W. SANDERSON & A. TABER. 2001. El Jaguar en el nuevo milenio: una evaluacion de su estado, detección de prioridades y recomendaciones para la conservación de los jaguares en America. México, D.F.: Universidad Nacional Autonoma de Mexico and Wildlife Conservation Society. 647 pp.
- MITTERMEIER, R.A., P.A. GIL, M. HOFFMAN, J. PILGRIN, T. BROOKS, C.G. MITTERMEIER, J. LAMOREUX, G.A.B. FONSECA. 2004. Hotspots revisited: Earth's biologically richest and most endangered terrestrial ecoregions. Mexico City: Agrupacion Sierra Madre. 392 pp.
- MORAES JR, E.A. 2012. The status of the Jaguar in the Cerrado. CAT news Special Issue 7: 25–28. [http://www.catsg.org/fileadmin/files/sharing/3.Conservation\\_Center/3.2\\_Status\\_Reports/Jaguar\\_Moraes\\_2012\\_The\\_status\\_of\\_the\\_jaguar\\_in\\_the\\_Cerrado.pdf](http://www.catsg.org/fileadmin/files/sharing/3.Conservation_Center/3.2_Status_Reports/Jaguar_Moraes_2012_The_status_of_the_jaguar_in_the_Cerrado.pdf)
- MORATO, R. G., B.M. BEISIEGEL, E.E. RAMALHO, C.B. CAMPOS & R.L.P. BOULHOSA. 2013. Avaliação do risco de extinção da onça pintada (*Panthera onca*) no Brasil. *Biodiversidade Brasileira* 3(1): 122–132. <http://www.icmbio.gov.br/revistaelectronica/index.php/BioBR/article/view/409/318>
- MYERS, N., R.A. MITTERMEIER, C.G. MITTERMEIER, G.A.B. FONSECA & J. KENT. 2000. Biodiversity hotspots for conservation priorities. *Nature* 403: 853–858. doi: [10.1038/35002501](https://doi.org/10.1038/35002501)
- NASCIMENTO, J.L. & I.B. CAMPOS. 2011. Atlas da fauna brasileira ameaçada de extinção em unidades de conservação federais. Brasília: Instituto Chico Mendes de Conservação da Biodiversidade. 276 pp.
- NÚÑES, R., B. MILLER & F. LINDZEY. 2000. Food habits of Jaguars and Pumas in Jalisco, Mexico. *Journal of Zoology* 252: 373–379. doi: [10.1111/j.1469-7998.2000.tb00632.x](https://doi.org/10.1111/j.1469-7998.2000.tb00632.x)
- ORTEGA-HUERTA, M.A. & K.E. MEDLEY. 1999. Landscape analysis of Jaguar (*Panthera onca*) habitat using sighting records in the Sierra de Tamaulipas, Mexico. *Environmental Conservation* 26 (4): 257–269. doi: [10.1017/S0376892999000387](https://doi.org/10.1017/S0376892999000387)
- PALMEIRA, F.B.I., P.G. CRAWSHAW JR., C.M. HADDAD, K.M.P.M.B. FERRAZ & L. VERDADE. 2008. Cattle depredation by Puma (*Puma concolor*) and Jaguar (*Panthera onca*) in central-western Brazil. *Biological Conservation* 141(1): 118–125. doi: [10.1016/j.biocon.2007.09.015](https://doi.org/10.1016/j.biocon.2007.09.015)
- PAVIOLI, A., C. DE ANGELO, K.M.P.M.B. FERRAZ, R.G. MORATO, J. MARTINEZ PARDO, A.C. SRBEK-ARAUJO, B.D.M. BEISIEGEL, F. LIMA, D. SANA, M. XAVIER DA SILVA, M.C. VELÁZQUEZ, L. CULLEN, P. CRAWSHAW JR, M.L.S.P. JORGE, P.M. GALETTI, M.S. DI BITETTI, R.C. DE PAULA, E. EIZIRIK, T.M. AIDE, P. CRUZ, M.L.L. PERILLI, A.S.M.C. SOUZA, V. QUIROGA, E. NAKANO, F. RAMÍREZ PINTO, S. FERNÁNDEZ, S. COSTA, E.A. MORAES JR & F. AZEVEDO. 2016. A biodiversity hotspot losing its top predator: the challenge of Jaguar conservation in the Atlantic Forest of South America. *Scientific Reports* 6: 37147. doi: [10.1038/srep37147](https://doi.org/10.1038/srep37147)
- PAVIOLI, A., C.D. DE ANGELO, Y.E. DE BLANCO & M.S. DI BITETTI. 2008. Jaguar *Panthera onca* population decline in the Upper Paraná Atlantic Forest of Argentina and Brazil. *Oryx Conservation Journal* 42: 554–561. doi: [10.1017/S0030605308000641](https://doi.org/10.1017/S0030605308000641)
- POLISAR, J., I. MAXIT, D. SCOGNAMILLO, L. FARRELL, M.E. SUNQUIST & J.F. EISENBERG. 2003. Jaguars, pumas, their prey base, and cattle ranching: ecological interpretations of a management problem. *Biological Conservation* 109: 297–310. doi: [10.1016/S0006-3207\(02\)00157-x](https://doi.org/10.1016/S0006-3207(02)00157-x)
- RABINOWITZ, A. & K. ZELLER. 2010. A range-wide model of landscape connectivity and conservation for the Jaguar, *Panthera onca*. *Biological Conservation* 143: 939–945. doi: [10.1016/j.biocon.2010.01.002](https://doi.org/10.1016/j.biocon.2010.01.002)
- REDFORD, K.H. & J.F. EISENBERG. 1992. Mammals of the Neotropics: the southern cone. Vol 2. Chicago: University of Chicago Press. 435 pp.
- REIS, N. R., A.L. PERACCHI, H. FANDINO-MARINO & V.J. ROCHA (orgs.). 2006. Mamíferos da Fazenda Monte Alegre – Paraná. Londrina: Editora da Universidade Estadual de Londrina. 176 pp.
- RIBEIRO M.C., J.P. METZGER, A.C. MARTENSEN, F.J. PONZONI & M.M. HIROTA. 2009. The Brazilian Atlantic Forest: How much is left, and how is the remaining forest distributed? Implications for conservation. *Biological Conservation* 142: 1141–1153. doi: [10.1016/j.biocon.2009.02.021](https://doi.org/10.1016/j.biocon.2009.02.021)
- RIPLE, J.W., J.A. ESTES, R.L. BESCHTA, C. CHRISTOPHER, E.G. WILMERS, M. RITCHIE, M. HEBBLEWHITE, J. BERGER, B. ELMHAGEN, M. LETNIC, M.P. NELSON, O.J. SCHMITZ, D.W. SMITH, A.D. WALLACH & A.J. WIRSING. 2014. Status and ecological effects of the world's largest carnivores. *Science* 343: 151–162. doi:
- RODRIGUES, E.H.G., I.M. MEDRI, W.M. TOMAS & G. MOURÃO. 2002. Revisão do conhecimento sobre a ocorrência e distribuição de mamíferos do Pantanal. Corumbá, MS: Embrapa Pantanal. 39 pp.
- RYALL, K.L. & L. FAHRIG. 2006. Response of predators to loss and fragmentation of prey habitat: a review of theory. *Ecology* 87: 1086–1093. doi: [10.1890/0012-9658\(2006\)87\[1086:ROPTLA\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2006)87[1086:ROPTLA]2.0.CO;2)

- SANDERSON, E.W., K.H. REDFORD, C.B. CHETKIEWICZ, R.A. MEDELIN, A.R. RABINOWITZ, J.G. ROBINSON & A.B. TABER. 2002. Planning to save species the Jaguar as a model. *Conservation Biology* 16(1): 58–71. doi: [10.1046/j.1523-1739.2002.00352.x](https://doi.org/10.1046/j.1523-1739.2002.00352.x)
- SANTOS-FILHO, M., C.A PERES, D.J. SILVA, T.M SANAIOTTI. 2012. Habitat patch and matrix effects on small-mammal persistence in Amazonian forest fragments. *Biodiversity and Conservation* 21: 1127–1147. doi: [10.1007/s10531-012-0248-8](https://doi.org/10.1007/s10531-012-0248-8)
- SCOGNAMILLO, D., I.E. MAXIT, M. SUNQUIST & J. POLISAR. 2003. Coexistence of Jaguar (*Panthera onca*) and Puma (*Puma concolor*) in a mosaic landscape in the Venezuelan llanos. *Journal of Zoology* 259: 269–279. doi: [10.1017/S0952836902003230](https://doi.org/10.1017/S0952836902003230)
- SEYMOUR, K.L. 1989. Mammalian species: *Panthera onca*. The American Society of Mammalogists 340: 1–9. <http://www.science.smith.edu/msi/pdf/i0076-3519-340-01-0001.pdf>
- SILVER, S.C., L.E. OSTRO, L.K. MARSH, L. MAFFEI, A.J. NOSS, M.J. KELLY, R.B. WALLACE, H. GÓMEZ & G. AYALA. 2004. The use of camera traps for estimating Jaguar *Panthera onca* abundance and density using capture/recapture analysis. *Oryx* 38: 148–154. doi: [10.1017/S0030605304000286](https://doi.org/10.1017/S0030605304000286)
- SOLLMANN R. 2010. Ecology and conservation of the Jaguar (*Panthera onca*) in the Cerrado grasslands of central Brazil [Dissertation]. Berlin: University of Berlin. 113 pp.
- SUNQUIST, M.E. & F.C. SUNQUIST, 2002. Wild cats of the world. Chicago: University of Chicago Press. 452 pp.
- TORRES, N.M., P. DE MARCO JR., J.A.F. DINIZ-FILHO & L. SILVEIRA. 2008. Jaguar distribution in Brazil: Past, present and future. *Cat News Special Issue* 4: 4–8. [http://www.jaguar.org.br/pdf/Publicacoes\\_4\\_Torres\\_et\\_al\\_2008\\_Jaguar\\_distribution\\_in\\_Brazil.pdf](http://www.jaguar.org.br/pdf/Publicacoes_4_Torres_et_al_2008_Jaguar_distribution_in_Brazil.pdf)
- TRAINOR, A.M., O.J. SCHMITZ, J. S IVAN, T.M. SHENK. 2014. Enhancing species distribution modeling by characterizing predator-prey interactions. *Ecological Applications* 24(1): 204–216. doi: [10.1890/13-0336.1](https://doi.org/10.1890/13-0336.1)
- UMETSU, F.; J.P. METZGER & R. PARDINI. 2008. Importance of estimating matrix quality for modeling species distribution in complex tropical landscapes: a test with Atlantic forest small mammals. *Ecography* 31: 359–370. doi: [10.1111/j.2008.0906-7590.05302.x](https://doi.org/10.1111/j.2008.0906-7590.05302.x)
- VILLARD, M.A & J.P. METZGER. 2014. Beyond the fragmentation debate: a conceptual model to predict when habitat configuration really matters. *Journal of Applied Ecology* 51: 309–318. doi: [10.1111/1365-2664.12190](https://doi.org/10.1111/1365-2664.12190)
- WHITE, P.C.L & P. LOWE. 2008. Wild mammals and the human food chain. *Mammal Review* 38: 117–122. doi: [10.1111/j.1365-2907.2008.00121.x](https://doi.org/10.1111/j.1365-2907.2008.00121.x)
- ZELLER, K. 2007. Jaguar in the new millennium data base update: the state of the Jaguar in 2006. New York: Wildlife Conservation Society-Jaguar Conservation Program. 77 pp.

**Received:** 17 January 2017**Accepted:** 9 March 2017**Academic Editor:** Guilherme S. T. Garbino